

Available online at http://www.ljerts.com

International Journal of Current Research in Life Sciences Fol. 07, No. 06, pp. 3220-2221, June, 2018



RESEARCH ARTICLE

BLOOD COAGULATION PROPERTIES OF KHANDU CHAKKA (EHRETIA LAEVIS) PLANT LEAVES

¹Diksha Bande and ^{2, 1}K<mark>shama Murarkar</mark>

¹Kamla Nehru Mahavidyalaya, India ²Assistant Professor Kamla Nehru Mahavidyalaya, India

Received 13th April, 2018; Accepted 06th May, 2018; Published 12th June, 2018

ABSTRACT

This study aims to investigate the potential of *Ehrtia laevis* as coagulant. To do this the dried powder of this plant was extracted with acctone and isopropanol using soxhlet assembly. These solvents were evaporated to make aqueous extract which was subjected to coagulation (prothrombin test) using plasma and blood sample of healthy individual. The blood coagulation time of prothrombin test of both the plasma and blood samples was reduced as compared to the control.

Key words: Ehrita laevis, coagulation time, Prothrombin time, extract

Copyright © 2018, Diksha Bande and Kshama Murarkar. This is an open access article distributed under the Creative Commons Attribution Decease, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Diksha Bande and Kshama Murarkar, 2018. "Blood Coagulation properties of Khandu chakka (Ehretia laevis) plant leaves" International Journal of Current Research in Life Sciences, 7, (06), 2220-2221

INTRODUCTION

Blood is a fluid in our body that helps to survive; transport all essential nutrients to body parts. Blood circulates through our body and delivers essential substances like oxygen and nutrients to the body's cells. There are four basic components that comprise human blood: plasma, red blood cells, white blood cells and platelets. It contains some factors that help to coagulate the blood when it flows outside the body. Coagulation is blood property that is evaluated in medical laboratories. Coagulation is an important mechanism of the blood to stop blood loss when damage occurs in the body Blood coagulants and anti-coagulants, 2008; Segen, 2004). There has been a number of literatures that studied coagulations (Maschirow et al., 2015; Luz et al., 2013; Vivas-Ruiz et al., 2013; Su et al., 2015; Menkiti et al., 2015; Merheb-Dini et al., 2012; Farhat et al., 2015). Romeo C. and Ongpoy J.R. (2016) studied coagulation and hemaggglutination properties of the crude extract derived from the leaves of Euphorbia Hirta L., Tridax procumbeans L., and Vernonia cinerea (L) Less (Romeo and Ongpoy, 2016). Essien et al. (1985) reported coagulant properties of root bark and root wood extracts of Fagara xanthoxyloides lam plant (Essien et al., 1985). Okuo and Ejimadu (2005) determined clotting time of blood samples drawn from eight patients using three variants of plant extracts (crude, aqueous and chloroform) of the leaf of Bryophyllum pinnatum. These extracts clotted the blood samples faster than the untreated blood samples, which ranged between 0.34-3.27 min (Okuo and Ejimadu, 2005). Edemeka and Oqwu (2000) effect of aqueous and methanol extracts of Ocimum gratissimum on the prothrombin time (PT)

and activated partial thromboplastin time (APTT) of plasma and factor VIII-deficient plasma. The biological activity suggests the extract contains unidentified constituents that promote blood coagulation (Edemeka and Oqwu (2000). In diabetic and high blood pressure patients' process of blood coagulation in wounds get delayed and blood flows continuously. This study was carried on blood coagulation property of Ehretia laevis (Khandu chakka). Ehretia laevis is a medicinal plant mainly used for wound healing, joint pain and minor fractures by local people with promising result. Thakre Rushikesh et al (2016) studied on ethano botanical properties of unexplored plant khandu chakka (Ehretia laevis roxb.), used on wound healing. Shailesh Dhenge and Kiran Khandare (2016) did a case study on the efficacy of local application of khandu chakka (Ehretia laevis roxb) ghrita in dushtavrana. The purpose of this study is to evaluate the coagulation properties of the Ehretia laevis.

MATERIALS AND METHODS

The plant material was collected from nearby jungle area of Nagpur. The samples were washed with distilled water and dried in sunlight to make powder form. The dried powdered material was extracted with soxhlet apparatus for 5-6 hours using acetone and isopropanol respectively. The extracts were collected and tested for coagulation activity (prothrombin test).

Collection of Blood sample for coagulation test

Blood sample from healthy individual having no disease and disorder was withdrawn from vein with the help of sterile syringe. It was placed in two separate vials containing trisodium citrate to prevent clotting of blood.

^{*}Corresponding author: Kshama Murarkar, Assistant Professor Kamla Nehru Mahavidyalaya, India.